Technical advice

Nemisia

Describtion:
Following early spring-flowering plants, nemesias are ready to market before the classic bedding and balcony plant season. Our series offer outstanding and unusual colours and very cold tolerant.

Potting:
Week 5 - 12
Pot 11 – 13 cm

Sales:
- Starts in march

Substrate:
- Potting-Substrate
- pH 5,5 – 6,5
- well aerated substrate with low salt content

Fertilizer:
Moderate nutrient need: with continuous fertigation at 0,05 % to 0,08 % or weekly fertigation 1 -2 times per week at 0,1 % bis 0,2 %.

Temperature:
Cold tolerant crop. Directly after potting daily average temperature 18°C, thereafter with DAT at 14°C.
Hardening at 10°C.

Light:
After roots established give full light during day

Flowers:
In march

Growth regulators:
Alar 1750 - 2500 ppm

Pinching:
Soft pinch after one week of growing.
Further pinch may be necessary depending on growth.

Diseases / pests:
Tospoviris, white fly, thrips, pythium, rizoctionia.

Spacing:
35 - 45 plants / sqm.

Elsner pac® Vertriebsgesellschaft mbH | Am Fiebig 14 | 01561 Thiendorf
Phone: +49 35248 3991-0 | E-Mail: info@pac-elsner.com

PASSION
AND COMPETENCE